

HHE UNITED STATES OF AMERICA

TO ALL TO VHOM THESE: PRESENTS SHALL COME: Hirginin Tech Intellectual Properties, Inc.

ILCCOS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT. THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID OPPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE NIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFFERING IT FOR SALE, OR REPRODUCING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE VERTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE E. OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT BY THE PLANT VARIETY PROTECTION ACT. IN THE UNITED STATES SEED OF THIS VARIETY (1) SO SOLD BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEO.)

WHEAT, COMMON

'3706'

In Testimonn Mucrest, I have hereunto set my hand and caused the seal of the Hant Hariety Frotestion Office to be affixed at the City of Washington, D.C. this twenty-ninth day of September, in the year two thousand and six.

Allast.

Range

Commissioner
Plant Variety Protection Office
A. h. h. h. M. l. v. C.

Agriculture

U.S. DEPARTMENT OF AGRICULTURE
AGRICULTURAL MARKETING SERVICE
SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE

The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995.

Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).

APPLICATION FOR PLANT VARIETY PROT (Instructions and information collection burden statement	ECTION CERTIFICAT	ΓΕ Application is n (7 U.S.C. 2421,	equired in order to determine if a plant van). Information is held confidential until cen	iety protection certificate is to be issued tificate is issued (7 U.S.C. 2426).
NAME OF OWNER Virginia Tech Intellectual Properties, inc.			TEMPORARY DESIGNATION OR EXPERIMENTAL NAME VA98W-706	3. VARIETY NAME 3706
 ADDRESS (Street and No., or R.F.D. No., City, State, and ZIF Virginia Tech Intellectual Properties, Inc. 1872 Pratt Dr., Suite 1625 	Code, and Country)		5. TELEPHONE (include area code 540-951-9378	PVPO NUMBER
Blacksburg, VA 24060			6. FAX (include area code) 540-951-5292	FILING DATE
IF THE OWNER NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZATION (corporation, partnership, association, etc.) Corporation	8. IF INCORPO STATE OF IN Virgin	ICORPÓRATION	9. DATE OF INCORPORATION June 20, 1985	August 15,2005
10. NAME AND ADDRESS OF OWNER REPRESENTATIVE (S) TO CARL A. Griffey Crop and Soil Environments Virginia Tech Blacksburg, VA 24061-0404	al Sciences	N. (First person listed w	ill receive all pápers)	FILING AND EXAMINATION FEES: \$ 3652.00 R DATE 8/15/2005 CERTIFICATION FEE: \$ 768.00
11. TELEPHONE (include area code) 12. FAX (include a 540-231-9789 540-231-343		E-MAIL Cgriffey@vt.edu	į į	DATE 4/2 (a/C)(a) CROP KIND (Common Name) neat, Common
 18. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMI reverse) a. X Exhibit A. Origin and Breeding History of the Variety b. X Exhibit B. Statement of Distinctness c. X Exhibit C. Objective Description of Variety d. X Exhibit D. Additional Description of the Variety (Option) e. X Exhibit E. Statement of the Basis of the Owner's Own f. X Voucher Sample (2,500 viable untreated seeds or, for verification that tissue culture will be deposited and ma repository) g. X Filing and Examination Fee (\$3,652), made payable to States" (Mail to the Plant Variety Protection Office) 	nal) Iership tuber propagated verieties, Intained in an approved public	20. DOES THE CONTROL OF T	YES (If "yes", answer items 20 and 21 below) DWNER SPECIFY THAT SEED OF THIS LIMITED AS TO NUMBER OF CLASSES CH CLASSES? X FOUNDATION DWNER SPECIFY THAT THE CLASSES ETO NUMBER OF GENERATIONS?	Veriety Protection Act) NO (If "no," go to item 22) X YES NO REGISTERED X CERTIFIED YES X NO REGISTERED CERTIFIED
22. HAS THE VARIETY (INCLUDING ANY HARVESTED MATERIAL) FROM THIS VARIETY BEEN SOLD, DISPOSED OF, TRANSFERI OTHER COUNTRIES? X YES IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALE, DISP FOR EACH COUNTRY AND THE CIRCUMSTANCES. (Please to	RED, OR USED IN THE U.S. OF	PROPERTY I	ETY OR ANY COMPONENT OF THE VAR RIGHT (PLANT BREEDER'S RIGHT OR F S COUNTRY, DATE OF FILING OR ISSUAN NUMBER. (Please use space indicated	PATENT)? X NO NOE AND ASSIGNED
24. The owners declare that a viable sample of basic seed of the variety for a tuber propagated variety a tissue culture will be deposited in The undersigned owner(s) is(are) the owner of this sexually reproduced in the entitled to protection under the provisions of Section 42 of the Owner(s) is (are) informed that false representation herein can jee	uced or tuber propagated plant ne Plant Variety Protection Act.	variety, and believe(s) th		
Muhay Met		SIGNATURE OF	OWNER	
NAME (Please print or type) (Michael J. Martin		NAME (Please pri	int or type)	•
CAPACITY OR TITLE Executive Vice President	DATE 5/25/05	CAPACITY OR TI	πE	DATE

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$2,450 (\$300 filing fee and \$2,150 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 500, NAL Building, 10301 Baltimore Avenue, Betsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$300 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291 Homepage: http://www.ams.usda.gov/science/pvp.htm

ITEM

18a. Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 21. See Section 83 of the Act for the Contents and Term of Plant Variety Protection.
- 22. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 5.5 of the Act for instructions on claiming the benefit of an earlier filing date.
- 21. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)

22. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

A limited amount of Certified seed of '3706' Wheat was sold in the U.S. A. for the first time in October 2004.

23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the variety names proposed by contacting: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center-East, Beltsville, MD 20705. Telephone: (301) 504-8089.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless if displays a valid OMB control number. The valid OMB control number for this collection of information is (0581-0055). The time required to complete this information is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data source gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and manital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964



18A. Exhibit A: Origin and Breeding History

Genealogy and Breeding Method.

Wheat cultivar 3706, previously tested as VA98W-706, was derived from the cross of VA91-54-350//FFR555W'/GA-Gore'. Parentage of VA91-54-350 is IN71761A4-31-5-48 //VA71-54-147 (CItr 17449)/ 'McNair 1813'. Wheat line IN71761A4-31-5-48 was developed by Purdue University and has the pedigree 'Benhur' (CItr 14054)/3/'Arthur' (CItr 14425)/'Knox' (CItr12798) type line/4/'Beau' (CItr17420)*2 /3/Arthur*2//'Riley' (CItr13702)/ 'Bulgaria 88' (PI 94407). The Knox type line has gene H5 for Hessian fly [Mayetiola destructor (Say)] resistance. The cross from which 3706 originated was completed in spring 1992, and the F1 generation was grown in the field at Warsaw, VA as a single 4ft headrow in 1993 to produce F2 seed. The population was advanced from the F2 to F4 generation using a modified bulk breeding method.

Population Advancement and Selection of the Variety.

Wheat spikes were selected from the population in each segregating generation (F₂-F₃) on the basis of absence of obvious disease, early maturity, short straw and desirable head shape and size. Selected spikes were threshed in bulk, and the seed was planted in 225ft² blocks in the fall of each year at Blacksburg and Warsaw, VA. Spikes selected from the F₄ bulk where threshed individually and planted in separate 4ft headrows at Warsaw, VA. Wheat cultivar 3706 was derived as a bulk of one of these F_{4:5} headrows selected in 1997, and was evaluated as VA98W-706 in non-replicated observation yield tests in 1998 as entry 706. It was evaluated as VA98W-706 in replicated preliminary yield tests in 1999 (Table 12) and in advanced yield tests in 2000 (Table 11). Wheat cultivar 3706 was tested as VA98W-706 in 2001 and 2002 under conventional (Tables 1-3) and no-tillage (Tables 4-6) management systems in the Virginia Official Variety Trial. It also was evaluated throughout the eastern soft red winter wheat region in the USDA-ARS Uniform Eastern Soft Red Winter Wheat Nurseries in 2002 (Table 8).

Multiplication and Purification.

Breeder seed of 3706 was derived as a bulk of 138 F_{8:9} individual headrows selected among 280 total rows evaluated for homogeneity and trueness of type in an isolation block at Warsaw, VA in 2001. During fall 2001, approximately 110 lbs of the 3706 Breeder seed was planted on 1 acre at the Virginia Foundation Seed Farm at Mt. Holly, VA and produced 78 bushels of Foundation Seed. During fall 2002, 30 acres of 3706 were planted and produced approximately 2000 bushels of Foundation seed that was available to seedsmen in fall 2003. While 3706 has remained stable and uniform in composition through the last three generations of self pollination, variants observed within the variety include up to 1% plants that are 6-8 inches taller in height, 0.5% plants with awned spikes that are similar in height to 6 inches taller, 0.1% plants having crooked or nodding spikes, and 0.01% plants having yellow green foliage and spike color.

'3706' Wheat

18B. Exhibit B: Novelty Statement

Wheat cultivar 3706 is uniquely different from all known wheat cultivars. It is most similar to its parent 'FFR 555W'. Wheat cultivar 3706 is resistant to powdery mildew (Blumeria graminis), while FFR555W is susceptible (See Table below). FFR555W possesses the gene Lr10 governing resistance to leaf rust (Puccinia triticina) and seedlings of FFR555W are resistant to leaf rust race TLGG (Virulence for genes Lr1, 2a, 2c, 3, 9, 11, 18) while seedlings of 3706 are susceptible to race TLGG. While the identity of Lr resistance genes of 3706 is not known, it consistently has expressed higher levels of resistance to leaf rust than FFR555W in field tests (See table below). Head emergence of 3706 is consistently earlier (2-4 days) than FFR555W. Test weight of 3706 also is consistently higher than that of FFR555W.

Comparison of 3706 with Parent FFR555W in 1999 – 2001 Virginia Replicated Variety Yield Trials

Variety	Grain	Test	Date	Plant		Powdery	Leaf
	Yield	Weight	Headed	Height	Lodging	Mildew	Rust
	Bu/Ac	Lb/Bu	Julian	Inches	0.2 - 10	0-9	0-9
<u>1999</u>	N=4	<u>N=4</u>	<u>N=2</u>	N=2	N=3	N=2	N=3
3706	86	60.1	124	33	0.3	0	1
FFR555W	72	58.3	128	35	0.4	6	4
L.s.d 0.05	5.8	0.5	1	1	0.6	1	1
2000	N=4	N=4	N=2	N=2	N=2	N=2	N=3
3706	90	58.2	120	34	1.0	1	2
FFR555W	70	55.5	124	35	0.2	6	7
L.s.d 0.05	7.3	0.9	3	1	0.9	2	1
2001	<u>N=7</u>	N=7	N=4	N=3	N=5	N=4	N=1
3706	81	58.5	123	29	2.0	2	0
FFR555W	76	56.2	125	32	1.3	6	4
L.s.d 0.05	4	1.2	1	2	1.1	1	1

Exhibit C (Wheat)

- OMB No. 9581-0955

REPRODUCE LOCALLY. Include form number and date on all reproductions.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 2.5 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, mental or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Breitle, large print, audiotape, etc.) should contact USDA7s TARGET Center at 202-720-2800 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or cell 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PLANT VARIETY PROTECTION OFFICE **BELTSVILLE, MD 20705**

OBJECTIVE DESCRIPTION OF VARIETY WHEAT (Triticum spp.)

NAME OF APPLICANT(S)	FOR OFFICIAL USE ONLY
Virginia Tech Intellectual Properties,	PVPG NUMBER
ADDRESS (Street and No. or RD No., City, State, and Zip Code)	200500325
Virginia Tech Intellectual Properties,	Inc. VARIETY NAME
1872 Pratt Dr., Ste. 1625	3706
Blacksburg, VA 24060	TEMPORARY OR EXPERIMENTAL DESIGNATION VA98W-706
minimum of 100 plants. Comparative data should be determined from varieties of used to determine plant colors; designate system used:	e number that describes the varietal character of this variety in the boxes below. 19 or less or 9 or less respectively. Data for quantitative plant characters should be based on a sentered in the same trial. Royal Horticultural Society or any recognized color standard may be swer all questions for your variety; lack of response may delay progress of your application.
1. KIND:	2. VERNALIZATION:
1 1=Common	2 1=Spring
2=Durum	2=Winter
3=Club 4=Other (SPECIFY) :	3=Other (SPECIFY):
3. COLEOPTILE ANTHOCYANIN:	4. JUVENILE PLANT GROWTH:
1 = Absent 2 = Present	1 = Prostrate 2 = Semi-erect 3 = Erect
5. PLANT COLOR (boot stage):	6. FLAG LEAF (boot stage):
3 1 = Yellow-Green	1 1 = Erect
2 = Green	2 = Recurved
3 = Blue-Green	1 = Not Twisted
	1 = Not 1 wisted 2 = Twisted
	1 = Wax Absent 2 = Wax Present
7. EAR EMERGENCE:	
1 2 0 Number of Days (Average)	
O O Number of Days Earlier Than	*
Same as McCo	ormick *
O 2 Number of Days Later Than 381	58 *
	* Relative to a PVPO-Approved Commercial Variety Grown in the Same Trial

				Exhibit C (Wheat
8. ANTHER COLOR: 1 = Yellow 2 = Purple			200500	
9. PLANT HEIGHT (from soil to top of head, 8 4 cm (Average)	excluding awns):			
o o cm Taller Than Same as	McCormick		*	• • • • • • • • • • • • • • • • • • •
0 8 cm Shorter Than	38158		*	
10. STEM: A. ANTHOCYANIN	D. INT	ERNODE		
1= Absent 2 = Present	1 5	1 = Hollow 2 = So Number of Nodes	emi-solid 3 = Solid	
B. WAXY BLOOM 1 = Absent 2 = Present	E. PE	OUNCLE 1 = Erect 2 = R	decurved 3 = Semi-e	rect
C. HAIRINESS	F. AUF	cm Length		
(last internode of rachis) 1 = Absent 2 = Present	1	Anthocyanin Hair Few		Present Present
11. HEAD (at Maturity): A. DENSITY				
1 = Lax 2 = Middense (Laxidense) 3 = Dense	2	RVATURE 1 = Erect 2 = Inclined 3 = Recurved		
B. SHAPE	D. AWI	NEDNESS	٠.	
1 = Tapering 2 = Strap 3 = Clavate 4 = Other (SPECIFY):	ر السيا خ	1 = Awnless 2 = Apically Awnletted 3 = Awnletted 4 = Awned		* • · · · · · · · · · · · · · · · · · ·
·	1		•	•

I. GERM SIZE

2

1 = Small

2 = Midsize 3 = Large

2

3 = Width Nearly as Wide as Kernel

1 = Depth 20% or less of Kernel 2 = Depth 35% or less of Kernel

3 = Depth 50% or less of Kernel

S&T-470-6 (07-02) designed by the Plant Varie	

Russian Aphid (Diuraphis noxia)

Cereal Leaf Beetle (Oulema melanopa)


Other (SPECIFY)

Other (SPECIFY)

1

0

	<u> </u>								Exb	ibit C (Wheat)
15.	INSECT: Continued	(0=Not Tested;	1=Susceptible;	2=Resis	tant;	3=Intermediate;	4=Toler	ant)		
		. 1	PLEASE SPECIFY	BIOTYF	E (whe	re needed)	200	e a	a z	9 8
0	Greenbug (Schize	iphis graminum)	•		Other	(SPECIFY)		o v	v J	
1	Aphids				Other	(SPECIFY)				
16.	ADDITIONAL INFO	RMATION ON A	NY ITEM ABOVE	OR GE	VERAL	COMMENTS	· · · · · · · · · · · · · · · · · · ·			



REFERENCE

Briggle, L.W. and L.P. Reitz. 1963. <u>Classification of Triticum Species and of Wheat Varieties Grown in the United States</u>. Technical Bulletin 1278. United States Department of Agriculture.

'3706' Wheat

18D. Exhibit D: Additional Description of '3706' Wheat.

Performance of 3706 (VA98W-706 in Tables) in comparison with other soft red winter wheat cultivars is presented in Tables 1-6 and 8. VA98W-706 is a high yielding, moderately early heading, short-stature, awnleted, soft red winter wheat with high test weight and good straw strength. Head emergence is 1 to 2 days later than 'Sisson' and 2 to 3 days earlier than 'Roane'. Average plant height of 3706 (31 inches) is similar to that of Sisson and 1 to 2 inches shorter than Roane. Straw strength (0.2=No lodging, 10=Completely lodged) of 3706 is good (range of 0.2 to 2.0 and mean of 1.2) and is better than that of Sisson (range of 1.5 to 4.3 and mean of 2.9). In Virginia's Official Variety Trials, average grain yields of 3706 in 2001 (81 bu/ac) and 2002 (85 bu/ac) under conventional tillage were significantly ($P \le 0.05$) higher than the test average (Tables 1-3). In no-till trials following maize conducted at Warsaw, Virginia, 3706 produced grain yields (89 bu/ac) similar to the test average in 2001, while grain yields of 3706 (91 bu/ac) were significantly higher than average in 2002 (Tables 4-6). Grain volume weight of 3706 (59.0 lb/bu) is high; it is 0.5 lb/bu lower than that of Roane and 1 lb/bu higher than that of Sisson.

Wheat cultivar 3706 was evaluated as VA98W-706 in the 2002 USDA-ARS Uniform Eastern Soft Red Winter Wheat Nursery (Tables 8), and ranked 5th among 44 entries in the region for grain yield (78 bu/ac), while the check variety Roane ranked 16th. In the same trials, wheat cultivar 3706 ranked 5th for grain volume weight (58.8 lb/bu). Based on data from four test sites in the Uniform Eastern SRW Wheat Nurseries, winter-survival of 3706 is good and similar to that of 'Caldwell' and 'Tribute'.

Reaction of 3706 (VA98W-706) to disease and insect pests has been evaluated in Virginia (Tables 1-6) and over the eastern region (Table 8). Wheat cultivar 3706 is resistant to powdery mildew (Blumeria graminis), and consistently had low scores (0-2 on a 0=Resistant to 9=Susceptible scale) in five states across the eastern region and Ontario, Canada. In seedling tests conducted at the USDA-ARS Cereal Disease Laboratory in St. Paul, MN, 3706 expressed resistance to 7 of 9 races of leaf rust (Puccinia triticina) and was resistant to 4 of 6 races of stem rust (Puccinia graminis). In field tests conducted over the eastern region, 3706 had leaf rust scores (0=Resistant to 9=Susceptible) ranging from 0 to 4 with an average score of 1.4 (Table 8). In tests conducted in Arkansas and Washington, 3706 expressed resistance to stripe rust (Puccinia striiformis). It has expressed high levels of resistance to soil borne mosaic virus and wheat spindle streak mosaic virus. Wheat cultivar 3706 is moderately resistant to leaf blotch (Septoria tritici), glume blotch (Stagonospora nodorum), and barley yellow dwarf virus. It is moderately susceptible to fusarium head blight (Fusarium graminearum). On the basis of seedling tests conduct by USDA-ARS at West Lafayette, IN, 3706 is susceptible to Hessian Fly [Mayetiola destructor (Say)] biotypes B, C, D, E, and L.

Milling and baking quality of 3706 (VA98W-706) is good and superior to that of Roane, Pioneer Brand '2580', 'Coker 9663', and 'USG 3209' (Tables 13-19). In comparisons with these four cultivars, 3706 yields more flour that is lower in water absorption and producers cookies of larger diameter. Flour protein concentration of 3706 (9.22%) is higher than that of FFR 555W (8.38%). On the basis of lactic acid retention capacity, flour of 3706 has stronger gluten strength (115%) than that of FFR 555W (102%). Therefore flour of 3706 likely will serve dual purposes in production of traditional pastry products as well as crackers.

Table 1. Two-year summary of performance of VA97W-24 and VA98W-706 in the Virginla Tech Wheat Test, 2001 and 2002 harvests.*

													ł
	٠	Test		Date						יפטאיייר	>	,00	
	Yield	Weight		Headed		Height		Lodaina		Mildew	× .	רכם קייני	
Line	(Bu/acre)	(Lb/bu))	Mar 31+)	<u>.</u>) E		(0.2-10)	. '	6-0		(6-0)	
	(13)	(13)		(8)		(9)		8		8		<u> </u>	
VA97W-24(RT)	84	58.0	,	č	+		. 4	C				•	•
VA98W-706(RT)	81	58.6	:	5 6		3 6	ا ۱	, k		. · ·		· (
TRIBUTE(RT)	+ 4	60.3	-	29	 	6 m	, ,	5 0		V C		ءاد	•
McCORMICK(RT)	+ 82	59.8	+	30		<u>,</u>	1	, c	,) c		> c	ŧ
SS550	84	57.5	ı	29	•	32		- c		ء د م) C	1 4
SISSON (RT)	84	57.8		27	- 5. - 1	ا ا	.1	3 6		1 C		۷ ۳	- -1
SS520	84	57.2		27	. 1	35	+	2.8		1 0		י כ	
USG 3209(RT)	82	57.3	`. J	28		30		2.8	٠.	1 0) e	1
PIONEER 26R24(B)	82	57.9	,	78 78	١,	32	+	24		1 6		2 6	- -
CENTURY II(D)	- 22	58.1		29	7,	33	+	3 1		7 7	+	o ←	+
AGS2000	75 -	58.0		27		32		. 4		٠,		- c	
FFR 518(RT)	74 -	56.8		27		<u>بع</u>	,			1 ~		o c	
PIONEER 26R38(B)	73 -	57.3	1	28		35	+	300		C	ı	<u>ر</u>	. 4
PIONEER 26R61(B)	- 69	59.2	+	58	•	8	÷	, & , &		ιn	+	4 C	۱ ۱
Toot Average (med 4)	Č	Ç	·		-								
ical Avelage (II-14)	<u>.</u>	58.4		သ		32		2.8				—	
C.V.	တ	<u>.</u>		4		4		54.3		63		89	
L.S.D. (0.05)	4	0.4		_		ς		1.0		\ 		} ~	

location-years on which data are based. A plus or minus sign indicates a performance significantly above or below the test average. * Varieties are ordered by descending yield averages. The number in parentheses below column headings indicates the number of

▼ Belgian Lodging Scale = Area x Intensity x 0.2. Area = 1-10, where 1 is wheat unaffected and 10 is entire

plot affected and Intensity = 1-5, where 1 is wheat standing upright and 5 is wheat totally flat.

▲ The 0-9 ratings indicate relative disease intensity where 0=none and 9=total plant infection.

Table 2. Summary of performance of VA97W-24 and VA98W-706 in the Virginia Tech Wheat Test, 2002 harvest.*

												Barley		•	
	. '									4		Yellow			
		Test		Date	1.7				ሏ	wdery	Leaf	Dwarf	Stripe	Early	
		Weight		Headed		-leigh		dging		lildew	Rust	Virus	Rust	Height	
Line		(Lb/bu)	_	Mar31+		Œ)		.2-10		(6-O)	(6 - 0)	(6 - 0)	6-0)	(inches)	
	(2)	(2)		(4)		(3)		(3)		(3)	(2)	(2)	(L)	£	
VA97W-24		58.1		28		36		.3		-	2	က	0	6.7	
VA98W-706		59.4		56		32		e.	•	'	0	က	0	5.5	
TRIBUTE		2.09	١.	26		32	•	7.		,	0	₹"	0	5.0 -	
SISSON		58.6		25		32		īĊ			4	7	8	7.3	
SS 550(R)	_	58.3		56		34		က.	•	1	က	-	7	7.0	
McCORMICK		0.09		56		32		.5	•	-	0	2	0	5.5	
USG 3209(RT)		57.9		24			_	+ 0.2	•	ı _	ည	~	0	8.0	
SS 520(R)		58.2		24		36		6.0		· -	4	က	2	8.7 +	
PIONEER 26R24(B)	_	58.8		25		35		6.0	``	~	က	2		6.3	
AGS2000(TV)		58.7		24		33		5.0		~	~	က	0	+ 0.6	
CENTURY II(D)		58.8		56		8 4			•	+	0	က	0	7.7	
PIONEER 26R38(B)		57.9		56		38		- 8. +		CI.	7	α	0	10.3 +	
SS 518(R)		58.0		24		32		÷		,	0	ന	0	+ 8.6	
PIONEER 26R61(B)		59.9	_	25		35	- i. l	1.4		3 . +	1	3	0	9.8	
										i N					
Average (n=65)	8	58.6		5 0		34	Ţ.	0.0	-	7	7	8	0	7.2	
C.V.	7	[4		ന	w	89.5	.,	7	43	88		11.4	
LSD (0.05)	4	0.4		_		·	-	0.7			•		1	1.3	

weight data from Blackstone. A plus or minus sign indicates a performance significantly above or below the * Varieties are ordered by descending statewide yield averages. These averages do not include yield or test test average.

The number in parentheses below column headings indicates the number of locations on which data are based Belgian Lodging Scale = Area X Intensity X 0.2. Area = 1-10, where 1 is wheat unaffected and 10 is entire plot

The 0-9 ratings indicate degree to which plant is affected, where 0=none and 9=total plant affected affected and Intensity = 1-5, where 1 is wheat standing upright and 5 is wheat totally flat.

**Wheat line having white seed color.

Early height (March 8) is an indication of the daylength sensitivity of a variety. Taller varieties began jointing in early March and thus were taller.

Table 3. Summary of performance of VA97W-24, VA98W-706, and VA98W-750 in the Virginia Tech Wheat Test, 2001 harvest.*

•							
		Test	Date		•	Powdery	Leaf
	Yield	Weight	Headed	Height	Lodging ♥	Mildew	Rust
Line	(Bu/acre)	(Lb/bu)	(Mar 31+)	(In)	(0.2-10)	(0 -9)	(0-9)
	(7)	(7)	(4)	(3)	(5)	(4)	(1)
VA97W-24(RT)	81 +	58.2	35 +	33 +	3.9	1 -	1
VA98W-706(RT)	81 +	58.5 ±	33	29 -	2.0 -	2 -	0 -
VA98W-750(RT)	79	57.1	34 +	29 -	3.2	1 -	0 -
TRIBUTE(RT)	87 +	60.3 +	33	30	2.8	0,-	0 -
USG 3209(RT)	84 +	57.6	33	- 28 -	3.3	2 -	1
PIONEER 26R24(B)	84 +	57.8	33	34 +	3.4	3	2 +
SS550	83 +	57.5	33	29 -	3.7	2 -	1
SISSON (RT)	83 +	57.5	31 -	30	4.3 +	2 -	1
SS520	83 +	56.9	32 -	33 +	4.0	3	3 +
McCORMICK(RT)	83 +	60.0 +	34 +	29 -	2.6	0 -	0 -
CENTURY II(D)	83 +	58.2	33	31	4.0	4 +	0 -
FFR 518(RT)	78	56.6	32 -	30	53+	1 -	0 -
PIONEER 26R38(B)	76	57.2	33	34 +	3.9	1 -	1
AGS2000	75	57.9	32 -	31	4.6 +	2 -	0 -
PIONEER 26R61(B)	72 -	59.1 +	34 +	33 +	2.0 -	3	0 -
						2	
Test Average (n=71)	77	57.3	33	31	3.1	3	. 1
C.V.	. 8	3.5	3	, 5	5.7	43	47
S.D. (0.05)	4	1.2	1 .	2	1.1	1	1

^{*} Varieties are ordered by descending statewide yield averages. A plus or minus sign indicates a performance significantly above or below the test average. The number in parentheses below column headings indicates the number of locations on which data are based.

[▼] Belgian Lodging Scale = Area x Intensity x 0.2. Area = 1-10, where 1 is wheat unaffected and 10 is entire plot affected and Intensity = 1-5, where 1 is wheat standing upright and 5 is wheat totally flat.

[◆] The 0-9 ratings indicate relative disease intensity where 0=none and 9=total plant infection.

Table 4. Two-year summary of performance of VA97W-24 and VA98W-706 in the Virginia Tech No-Till Wheat Test at Warsaw, 2001 and 2002 harvests.*

Line	 Yield (Bu/ad		Tes Weig (Lb/b (2)	ht	Date Head (Mar 3 (2)	ed	Heig (In) (2)		Lodgin (0.2-1	_	Powd Milde (0-9 (2)	ew
VA97W-24	102	+	56.1		30	+	32	+	2.0		1	
VA98W-706	90		57.0		29		26	-	0.2		0	
TRIBUTE	99	+	59.4	+	29		28		0.4		0	-
SS550	 95	4	56.2		29		27		1.5		. 1	
SISSON	95		55.9		27	-	26		0.6		1	
SS520	 94		55.6	-	27	-	30	+	0.3		0	-
McCORMICK	93	4	58.4	+	30	+	27	-	3.4		0	-
PIONEER 26R24	 90		56.1		30	+	28		1.9		1	
USG 3209	88		55.7	-	29		26	-	1.2		1	
CENTURY II	 . 84		56.5		30	.+	28		1.1		2	+
FFR 518	83		55.4		30	+	27		4.4	+	0	.=-
PIONEER 26R38	 - 80	-	55.1	-	31	+	29	+	1.1		1	
AGS2000	79	-	56.6		28	-	28		3.2		1	
PIONEER 26R61	 75	-	57.6	+	30	+	30	+	0.2		2	+
Test Average (n=14)	89	.3	56.5		29	٠.	28		1.5		1	j
C.V.	8		1.4		3	- A - 2	. 4		103.8		75	
L.S.D. (0.05)	7	•	0.8		1		1		2.3		1	

^{*} Varieties are ordered by descending yield averages. The number in parentheses under column headings indicates the number of years upon which data are based. A plus or minus sign indicates a performance significantly above or below the test average.

[▶] Belgian Lodging Scale = Area x Intensity x 0.2. Area = 1-10, where 1 is wheat unaffected and 10 is entire plot affected and Intensity = 1-5, where 1 is wheat standing upright and 5 is wheat totally flat.

[♠] The 0-9 ratings indicate relative disease intensity where 0=none and 9=total plant infection.

Table 5. Summary of performance of VA97W-24 and VA98W-706 in the Virginia Tech No-Till Wheat Test at Warsaw, 2002 harvest.*

		Test			Date	ē.			Powdery	Leaf	Freeze	T
	Yield	Weight		Lodging	Headed	ded	Height	٠ .	Mildew	Rust	Damage	
9	(0(0)	(L. 1)		6			, ;		one rep	:	one rep	
רוום	(bu/acre)	(na/an)		(0.2-10)	(Mar31+)	31+)	(III)		(0-0)	(6-0)	(1-5)	
VA97W-24	10e +	54.2		2.0	22	+	31	+	2	2	2	
VA98W-706	91	54.2		0.2	20		28	ş	0	ß	\ (\	
TRIBUTE	101 +	57.3	+	0.4	20		29	١.	0	2	-	
McCORMICK	+ 86	56.1	+	3.4	22	+	30			က '	. ve-	
SS 550(R)	+ 96	53.2		7.5	20		30		-	+	. ო	7
SS 520(R)	91 +	52.8		0.3	20		31	+	7	ဖ	0	
SISSON	+ 06	52.3	,	9.0	18		28	•	m	+	۱ ۵	
PIONEER 26R24(B)	+ 68	53.2		1.9	20	. 1	30		က	+ &	ı 🛚	
CENTURY II(D)	84	53.6		<u>-</u> :	20		99		က	4	4	
USG 3209(RT)	80	52.7		1.2	20		27	٠.	4	Ŋ	4	
SS 518(R)	9/	52.3	1	4.4	20	•	28	1	0	رى د	4	
AGS2000(TV)	73 -	54.0		3.2	17	•	29	. 1	ო	4	4	٠.,
PIONEER 26R38(B)	- 02	51.3		- -	2		32	+	•	ເນ	4	
PIONEER 26R61(B)	- 65	55.6	+	0.2	2		31	+	7	က	4	
4								İ				
Location Average (n=65)	82	53.6	٠	-	7		ဓ္က		် က	ιΩ	24	
LSD (0.10)	7	*		4	<u> </u>		~		1	7	1	
c.v.	8	1.8		112.0	Ω		4		1	35	l	

* Varieties are ordered by descending yield averages. A plus or minus sign indicates a performance significantly above Belgian Lodging Scale = Area X Intensity X 0.2. Area = 1-10, where 1 is wheat unaffected and 10 is entire The 0-9 ratings indicate degree to which plant is affected, where 0=none and 9=total plant affected. or below the test average. There is only one rep of data for powdery mildew and freeze damage. plot affected and Intensity = 1-5, where 1 is wheat standing upright and 5 is wheat totally flat. The freeze damage ratings 1-5 are 1=no damage and 5=all early tillers killed.

Table 6. Summary of performance of VA97W-24, VA98W-706, and VA98W230 f) 5 () () 3 2 5 the Virginia Tech No-till Wheat Test at Warsaw, 2001 harvest.*

Line	Yiel (Bu/ad		Test Weight (Lb/bu)	Dat Head (Mar 3	ded	Heig (In		Powo Mild (0-	ew
VA97W-24(RT)	99	+	57.9	38	+	33	- , +	0	-
VA98W-706(RT) VA98W-750(RT)	89 89		59.7 58.8	35 38	+	25 24	-	0	-
SISSON (RT)	100	+	59.8	33		25	-	- 1	
USG 3209(RT)	97	+	58.2	36		25		0	
SS520	97	+	58.4	33	-	29	+	0	
TRIBUTE	97	+	61.1	35	-	27		0.	
SS550	94		59.3	35	-	24	-	0	-
PIONEER 26R38(B)	90		59.3	37	+	28	+	0	~
FFR 518(RT)	> 90		58.3	37	+	26		0	-
PIONEER 26R24(B)	90		58.7	37	+	27		2	+
CENTURY II(D)	88		59.3	37	+	27		2	+
McCORMICK	88	٠	60.6	37	+	25	, -	0	
PIONEER 26R61(B)	85		59.5	38	+	31	+	3	+
AGS2000	85		58.4	39	+ .	26	-	0	
Test Average (n=71)	88		58.3	36		27	:	1	
L.S.D. (0.05)	9		5.9	1	4 1	1		1	
C.V.	7		7.3	3		4		68	

^{*} Varieties are ordered by descending yield averages. A plus or minus sign indicates a performance significantly above or below the test average.

[•] The 0-9 ratings indicate relative disease intensity where 0=none and 9=total plant infection.

	Black	Blacksburg	Mean Y	rield	Mean T	Mean Test Wf.	Mean Head Date	d Date	Mean Plant Ht	Mean
		VA	All Lo	SOO	AIII	All Locs	All Locs	SS	Alllocs	16 l oce
Line	bu/ac	rank	bu/ac	rank	ng/sql	rank	Julian	rank	Inches	6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6
/A98W-706	72.2	27	72.0	7	58.8	5	130.4	4		23
Caldwell	64.3	39	58.5	43	56.6	32	135.0	34		1 c
Foster	78.6	Φ	9.99	30	57.3	24	134 7	S &	36.0	., c
Patton	75.4	18	70.6	10	57.6	G.	131.4	} «		2.5
Roane	82.6	7	69.4	4	59.8	2	135.1	, K	2 PK	7 C
Tribute (VA98W-593)	90.8	က	72.9	4	59.9	i -	131.9	.) c

			Winter Kill/Fre	reeze Damage			Powdery Mildew	Leaf Rust	Stem Rust
	Bay	Manhattan 7.5	Columbia	Columbia	Smithville	Same Brokens			1
	freeze (1.	2	2	freeze injury	survival	Survival	e Locs	/ Locs LA MN	
Line	6	winter dmg. survival (%)	survival (%)	(%)	(%)	Clar 1 7	6-0	(6-0)	
VA98W-706	1.3	3.0	91.0	1.3	100.0	0.88	1.2	0	S 60MS-S
Caldwell	1.3	4.0	87.0	2.0	98.0	88.7	4	50 MS	
Foster	0.	3.0	93.0	17	98.5	93.3	3.3	2.0 15 MS 40MR-MS	
Patton	1.7	2.0	93.0	1.3	100.0	95.3	တ	0	
Roane	2.0	4.0	0.68	2.7	95.0	97.3	1.7	7 MR	40MS-S
Tribute (VA98W-593)	10	3.0	91.0	1.0	100.0	0.7	0.4	1.5 0 20MR-MS	

	Stripe		e de la companya de l						Spindle	Soil Borne
	Rust	Ś	Septoria	ᆵ	Fusarium Head Blight (Scab)	Blight (Scab		BYDV	Streak	Mosaic
		Septoria tritici	Stagonospora nodorum	Urbana	Urbana	Urbana			clay Co.	
	2 Locs	5 Locs	Lexington, KY	4	4	1	2 Locs	5 Locs	AR	-
Line	%	6-0	-0-5	hcid. (%)	Sev. (%)	-0) (00) 100)	6-6	6- 0		6-O
VA98W-706	5.0	3.5	က	95.0	60.0	57.0	3.8	3.7	1.0	2.5
Caldwell	51.7	5.3	4	86.7	39.5	34.2	2.8	3.9	3.7	0.9
-oster	70.0	3.7	2	80.0	23.3	18.9	2.3	2.0	4.3	ر ئ
Patton	75.0	4.6	2	76.7	28.6	22.3	2.0	3.1	0	1.0
Roane	28.7	3.5	က	75.0	27.4	20.6	1.5	2.2	3.7	5.0
Fribute (VA98W-593)	27.4	2.5	က	85.0	33.3	27.9	G	er.	37	7.2

Table 11. Summary of performance of VA98W-706 in the 1999-2000 Advance Wheat Yield Test in Virginia over 4 locations. The numbers below each column heading indicate the number of locations upon which data are based.

		Percent		Heading	••					Plant Height	Juvenile Plant	
	Viold	of Test	Test	Date	Plant		Powdery	Leaf		o u	Growth	
Line	(bu/a)	(Yield)	(lbs/bu)	(Marcn 31+)	Height (in)	$(0.2-10)^1$	$(0.9)^2$	Rust (0-9)	BYDV	3/24/00 (in) ³	Habit	
	4	4	4	2	2	2	2		,	(III) -	(6-5)	
VA98W-706	6'68	107.0	58.2	30	34	10		,	1 (4	- =	-	
ROANE	82.3	0.86	58.7	33	34	7	, (r	1 4	, c	= =	→	
COKER 9663	81.1	96.5	58.0	30	40	2.1	, 4	· -	۱ ر	1 1	> c	
FFR 555W	70.4	83.8	55.5	34	35	0.2	ي .	- 1	, 1 (4	3 2	7	
PION 2580	9.98	103.1	56.2	53	38	0.5) (r	- и	, ,	<u> </u>	٦ ,	
PION26R24	89.5	106.5	57.2	30	38) (T	4	4 C	7 [4	
AGS 2000	88.1	104.9	57.5	28	36	4.	2	-	1 6	7 P	۰ ۲۰	
VA98W-750	91.0	108.3	57.0	33	35	0.3			۱ ر	<u> </u>	, c	
GRAND MEAN	84.3		56.9	31	36	1:0	2	۱۳	1	5 2	7 -	
CV	12.3		2.2	13	т	106.1	83	49	× 48	} _	74	
ISD	7.3		6.0	m	-	6.0	7		5		t ₇ 0	
											1	

¹ Belgian lodging scale = area x intensity x 0.2. Area is rated on a scale from 1 (plot unaffected) to 10 (entire plot affected). Intensity is rated on a scale from 1 (plants standing upright) to 5 (plants lying totally flat on the ground).

² All 0-9 ratings indicate relative disease severity: 0 = no disease present; 9 = total plant infection.

³ This height measurement serves as an estimate of spring growth habit.

⁴ Juvenile Plant Growth Habit: 0 = prostrate; 5 = very erect.

Virginia and Plymouth, North Carolina. The number below each column heading indicates the number of locations upon which Table 12. Summary of performance of VA98W-706 in the 1998-99 Preliminary Wheat Test at Blacksburg, Warsaw, and Painter, data are based.

		Test	Heading			Winter	Powdor			
Line	Yield (bu/a)	Weight	Date (Inlian)	Height	Lodging	Kill	Mildew (0.	Ę,	Septoria	BYDV
	4	4	2	(mm)	(0.2-10)	(6-2)	ع ا	(6-0)	(6-0)	(6-0)
1 1 1 0 0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1					,	7	7	ņ		7
VA98W-706	85.5	60.1	124	33	0.3	2	0	-	-	٠
Madison	77.7	57.4	122	37	0.8	8	2	• 🔻	- «	1 -
Agripro Patton	75.6	58.2	124	36	0.5	.0	۱۸	٠ -) [٦ ,
Roane	85.8	8.09	128	34	13	· C	í v-		⊣ <u>C</u>	ے ن ۔
Pocahontas	77.8	59.3	122	35	6.0	2 0		ዞ <u>፤ር</u>	4 -	- c
Coker 9663	75.6	59.3	123	38	8.0	l et	ı 4) (٦ -) t
Pioneer 2580	90.1	58.4	123	36	0.7) en		T V	- ا	٦ ,
FFR 555W	71.6	58.3	128	35	0.4	, , 4	ועכ	ዞ 뒥	1 +	4 c
VA98W-750	83.5	59.0	126	34	0.2	· C) [א כר	٦, ٣	4 c
VA98W-817	77.4	67.5	122	37	0.4	, 1) (-	· ·	1 4
Test Mean $(n=93)$	80.4	58.8	125	34	9.0	2		2 2	202	* ~
LSD (0.05)	5.8	0.5	, - 1		9.0	.	₩.	-	0.5	٠.

¹ Belgian Lodging Scale=area x intensity x 0.2. Area is rated on a scale from 1 (plot unaffected) to 10 (entire plot affected). Intensity is rated on a scale from 1 (plants standing upright) to 5 (plants lying totally flat on the ground).

² All 0-9 ratings indicated relative disease severity: 0 = no disease present; 9 = total plant infection.

Table 13. Milling and baking quality of VA98W-706 in the 2001-02 Uniform Eastern Soff Red Winter Wheat Nursery

	Mill Score	IIII Score Bake Score	Soft Equiv.	Ē	our Yield Flour Protein Gluten Streng	Gluten Strength	Water Abearh	Cookio Diam	
		;	%	•	%	(alter point and)	20000	3	
VA98W-706	92.7 C	88 5 D	19 E**	74 E	200	lactic acid let II)	%	동	
2 - 1 - 1 - 1 - 1 - C)	2	0.0	0, -	9.82	0.00	£7.8	47 04*	٠
Caldwell (standard)	100.0 A	100.0 A	50	ν 1.7	0 00	 	0.10	- S	
Foster	104 & A) (0.00	6.1.1	60.3	18.2	
; i	0.00	94.3 G	23.50	0.4/	9.44	ر در در	7.7	7 0 7	
Patton	90.8 C	82.8 E	53.3*	70 A*	0 83	5 6	1.70		•
Roane	96.6.0	л П	* 4 7 *	† *	0.00	94.0	61.9	17.82*	
MoCormick	2 0	- 1	 	900	9.58	117.1	65.4**	16.78**	
INCCOLLINGS.	91.0 C	73.9 F	54.8	**0.07	9.73	777	*0 00	2 1	
					5	T		**/V/L	-

*One standard deviation below the check variety
**Two standard deviations below the check variety

Mill Score E	Mill Score	Bake Score	Flour Yield	Soft Equiv.	Flour Protein	ore Bake Score Flour Yield Soft Equiv. Flour Protein Gluten Strength Water Absorb	Water Absorb
77.00011			%	%	%	(lactic acid ref'n)	%
VA98W-750	82.5 E	96.9 B	70.8**	57 98	777	00 75	2
VA98W-706	0,00	0.4	110	5	11.	07.70	57.4**
	3.5	3	74.0	48.08*	9,13	70 11	6 6
VA97W-24	91.8 O	94.8 C	×* 2 8 2 2 4 **	52 79	100	- 00	0.00
)	- -	07.70	0.00	/3.29	55.7*
roster (standard)	100.0 A	100.0 A	76.3	52.73	8 34	74.40	, I
Roane	מ 7 אמ	00.4	1	i	t o	2.4	22.7
	5	08 D	71.(~	53.60	8.06	85.30	EO 7**
Coker 9663	87.9 D	83.3 T	70 K**	*00 47		0.00	7.60
0000) (1	5.4	00.74	0/0	79.62	58 0**
FION 258U	88.2 D	84.8 E	72.6**	47 90*	α π	40.07	1 ()

*One standard deviation below the check variety

^{**}Two standard deviations below the check variety

Table 15. Milling and baking quality of VA98W-706 in the 2000-01 Virginia State Wheat Nursery.

	Mill Score	Mill Score Bake Score	Flour Yield	Soft Equiv.	Flour Protein	Gluten Strenath	Water Absorb	Cookie Diam
1100011			%	%	%	(lactic acid ref'n)	%	COORIE DIAIII,
VA38VV-/ U5	102.4 A	103.5 A	74.2	52.6	8.26	1160	200	C
VA97W-24	102.4 A	104 0 A	73.7	F: 0.4	1 (0.01	22.2	18.28
Dionage 28024	•	C + 0 + 0 + 0 + 0 + 0 + 0 + 0 + 0 + 0 +	- i	00.	8/./	86.0	57.2	18.20
רוטווככו בטרב4		100.0 A	72.3	56.7	7.79	114.3	9 04	7 1 1 1 1 1 1 1
Massey	100.1 A	105.2 A	727	54.3	α α	- *	50.00	8/:/1
Jackson	100.8 A	107.5 A	72.4	9 0	1 0	0.01	96.9	18.17
Roana	05.1 D	20.00	1.7.1	0.60	CS: /	122.6	58.9	18.25
Sign of Control of Con		IUZ.O A	/0./	58.2	7.33	110.8	60.5	18 21
OISSON	96.3 B	80.7 E	72.3	50.9	7.56	8 70	* C	5.0
USG 3209	86.5 D	85.3 D	**6 02	48 O##	7 20	0 7	00.00	17.34
Century II	105 B A	1 4 6 6	1 5		00.7		633	18.00
		£ 0.00.	13.1	5/.1	7.88	102.6	59.6	17 87
AGS 2000	104.1 A	105.8 A	74.7	55.0	7 64	105.0	9 6	0 0
Coker 9663	919 C	96 D B	74.0*	* 7 7 1	1 - 1	0.001	90.9	18.85
Coker 0835	104 6	2000	- 1	4	1.37	111.4	57.6	17.73
0.000	2 6	100.0 A	(3.1	59.4	7.29	95.7	59.7	18 49
Pioneer 2580	91.9 C	103.2 A	* 6.02	53.7	7.29	109 7	57.0	7 0
Pioneer 2684	97.0 B	104.3 A	72	72 73 74	000		2.10	08.7
Pioneer 2643	90 3 8	108.8 A	604	2 1	1 0.1	104.1	25.7	18.40
	0.00	C	72.3	22.8	1.72	126.4	56.7	18.53
Ploneer 25K45	102.7 A	104.4 A	75.5	53.5*	7.29	113.4		5 5
FFR555W	102.5 A	105.7 A	73.4	54.8	7.70	104 1	7.17	0.60
							33.	18.72

*One standard deviation below the check variety
**Two standard deviations below the check variety

23

Table 16. Milling and baking quality of VA98W-706 in the 2000-01 Virginia Elite Wheat Nursery.

		WIII SCOIE DAKE SCOIE	Flour Yield	Soft Equiv.	Flour Protein	Gluten Strength	Water Absorb	Cookie Diam
			%	*	%	(lactic acid ratin)	/0	
VA98W-750	85.9 D	93.8.0	80 2**	E2 4	40.704	ייייטיוט מסומ ובר וו)	-0/	CIII
1/4 CO14/ 100)	9	1.70		83.4	بر م	17.10
VASSW-/UD	95.6 B	101.5 A	72 A*	40.0	600		<u>.</u>	01.7
	1 1 1 1		į	10.0	20.0	114.2	48 2	47.44
rrkopovy (standard)	100.0 A	100.0 A	73.3	50.0	740		1	‡ . :
, COOC - COOC		0:00)	000	40.0	103.2	47.8	18.00
Coker 9003	89.5 D	0 0 0 0 0 0	**0 1/	17 2*	200	0 0 0	?	0.33
100 to 0)	į -	?	2.0	112.3	40 G*	46.00
ration	93.1 C	87.6 D	71.2**	500	10.04	000		76.01
Dog C	.0		į	7.70	5.0	89.0	513**	16.86
יסמונם	<u>ال</u> الم	83.8E	* *002	55.0	0 10	0.00		22.
1100 2200	£ 0	(ò	2.0	7.001	20.5	16.48**
920 9509	85.3 D	29.5 F	**. 69	400	10.19	4006	1) (

*One standard deviation below the check variety **Two standard deviations below the check variety

Table 17. Milling and baking quality of VA98W-706 in the 1999-2000 Virginia Adva

	Mill Soor	Mill Score Bath Start Flagger			1	ulacily.	
		Dane Score	riour Yield	Soft Equiv.	Flour Protein	Gluten Strength	Water Absorb
VA98W-750	89.6 D	92.4 C	70 6**	70 52	%	(lactic acid ret'n)	%
VA98W-708		ie	7,0	25.87	က က	75.9	59.3*
	20.0	ည် (၁၈	74.8	49.43*	9 27	0) ⁽
FFR555W (standard)	100.0 A	100 0 A	75.0	90 7 11	4 0	9.00	57.3
Roane	ARA D	7 0 00	2.0.0	04.20	g.65	77.9	54.9
0000	1 1 0	9.00 j	72.5	55.52	8.37	78.8	#B 07
JUNEI 8003	92.5 C	87.8 D	73.5**	50.28*	α α	7	0 1
Pion 2580	87.5 D	96 4 B	74 0**	70 00		0.17	59.3*
Pion 26R24	0 00		- i	90.00	7.23	72.8	58.1*
	32.30	99.0 B	(3.7"	58.05	8.02	848	*0 04
						ָ כֿי כֿי	0000

*One standard deviation below the check variety

Table 18. Milling and baking quality of VA98W-706 in the 1998-99 Virginia Preliminary Wheat Nursery.

	Mill Score	Bake Score	Elone Viola	- L 13-0				٠.	
				Sort Equiv.	Flour Protein	Gluten Strength	Water Absorb	Cookie Diam.	
7700007			/0	%	%	(lactic acid ref'n)	%		
VA86VV-75U	93.4 C	84.4 E	70.07	57.0	V 0	4000	9/	ES	
VA98W.706	V 0 00	100) i	50.0	106.0	20.2	18 13	
	Į,	1.0	8.7	51.2	00 20 20 20	1427	40 01		
VA98W-817	79.9 F	61.7 F	R7 0**	*C (C)	0	t 5	20.00	17.78**	
Madison (etopologia)	4000		5	43.7	Ø. 58	139.3	50.2**	17 47**	
iviacioni (stalicalo)	40.00 H	100.0 A	72.6	53.1	8 29	4010	1 1	- t	
Patton	99.0 B	90 0 0	2	7 7 3	1 0	6.101	04.Z	18.36	
0.000) () : i :	7	4.40	/0./	ත <u>්</u> ත	57.0**	18 20	
Noarie	91.1 C	61.4 F	£4. 20.7**	54 4	7.52	1047) i	0.00	
Coker 9663	89.3 D	50.5 F	**V UZ	***	20.0	7.001	52.4**	17.48**	
Diopost 2500	1 1	- I	+	12	8.07	108.2	÷00.6**	16 98**	
Liolieel 2000	8/.1D	58.4 F	\$ 1	53.6	7 23	0 707	; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ;	9	
FFR555W	105.9 A	01.8	72.8	1 (- 1 0 1	0.101	50.6**	17.14**	
		9	0.5	04.7	28	00 00 00	* 20 20 20 20 20 20 20 20 20 20 20 20 20	10.07	
			!			1			

*One standard deviation below the check variety **Two standard deviations below the check variety

Table 19. Milling and baking quality of VA98W-706 in the 1997-98 Virginia Wheat Observation Nurs

	Will Score	Bake Score	Flour Yield	Soft Equiv.	Flour Protein	Water Absorb
			%	%	%	2
VA98W-750	92 8 C	02.7 0	10.04		0/	0/
) Pii:		12.2	54.74	10.70	56 7**
VA98W-706	100.9 A	92.6 C	7.47	*66 07	7	Ö
Madioon (propage)	0007			43.43	10.46	53.7
iviacisori (standard)	100.0 A	100.0 A	74.4	52 RA	80 C	
FFR555W	40004	000	· 1	10.1	9.00	1.76
	C 0.001	30.0 B	/4./	53,55	8 45	* 7 7 4
Jackson	סקים	0.4.2.0	10 04) (P S	4.40
	0.00	a c . / a	73.0*	56.29	71.6	15 O.*
Pioneer 2580	0000	00 0 B	10 0**		· (7.00
) ii	2	1.7.7	55.23	00 00 00 00 00 00 00 00 00 00 00 00 00	35 XX
Pioneer 2552	100.9 A	9200	777	60.44	, ,	9.
07.00		0	į	28.10	9.64	56 O*
Florieer 2543	95.9 B	98.3 B	73.1*	55.07	Č) (
Dionagr 2804		0 707	· · ·	50.00	- O.9	55.3*
1 2011051	90.4	101.9 A	71.5**	29 77	0 73	200
Coker 0803	000	1		;	5	93.9
2000 1000	30.0E	20.00	/3.2*	56.12	8 04	# C 14
Coker 9835	943 C	96 A B	** 1 0 1		† ()))	0.00
	5 5:: 5	00.00 D	1.2.1	51.88	7.86	57 O**

*One standard deviation below the check variety
**Two standard deviations below the check variety

REPRODUCE LOCALLY. Include form number and edition date on a	Il reproductions.	FORM APPROVED - OMB No. 0581-005			
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E	Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). The information is held				
STATEMENT OF THE BASIS OF OWNERSHIP	confidential until the certificate is issu	ued (7 U.S.C. 2420).			
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3. VARIETY NAME			
Virginia Tech Intellectual Properties, Inc.	OR EXPERIMENTAL NUMBER	3706			
	VA98W-706	0.00			
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (Include area code)	6. FAX (Include area code)			
1872 Pratt Drive	1	, ,			
Suite 1625	540-951-9374	540-951-5292			
Blacksburg, VA 24060	7. PVPO NUMBER				
)			
8. Does the applicant own all rights to the variety? Mark an "X" in the	4	200500325			
9. Is the applicant (individual or company) a U.S. national or a U.S. I					
10. Is the applicant the original owner?	X NO If no, please answer on	e of the following:			
a. If the original rights to variety were owned by individual(s), is (are) the original owner(s) a U.S. National(s)? YES NO If no, give name of country					
b. If the original rights to variety were owned by a company(ies) X YES 11. Additional explanation on ownership (If needed, use the reverse	NO If no, give name of coul				
Original owner Virginia Polytechnic Institute and State Universit Properties, Inc. (See Attached document).	•	owner Virginia Tech Intellectual			
PLEASE NOTE:	· · · · · · · · · · · · · · · · · · ·				
Plant variety protection can only be afforded to the owners (not licens	sees) who meet the following criteria:				
If the rights to the variety are owned by the original breeder, that p national of a country which affords similar protection to nationals or	person must be a U.S. national, national of the U.S. for the same genus and spec	of a UPOV member country, or eles.			
If the rights to the variety are owned by the company which employ nationals of a UPOV member country, or owned by nationals of a genus and species.	yed the original breeder(s), the compan country which affords similar protection	y must be U.S. based, owned by to nationals of the U.S. for the same			
3. If the applicant is an owner who is not the original owner, both the	original owner and the applicant must n	neet one of the above criteria.			
The original breeder/owner may be the individual or company who di Act for definitions.	irected the final breeding. See Section	41(a)(2) of the Plant Variety Protection			
According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, control number. The valid OMB control number for this information collection is 0581-0055. including the time for reviewing the instructions, searching existing data sources, gathering	. The time required to complete this information colle	ction is estimated to average 0.1 hour per response,			
The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and a marifal or family status, political beliefs, parental status, or protected genetic information. (In communication of program information (Braille, large print, audiotape, etc.) should contact U	Not all prohibited bases apply to all programs.) Perso	ons with disabilities who require alternative means for			
To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 328- 720-5984 (voice and TDD), USDA is an equal opportunity provide and employer.	W, Whitten Building, 14th and Independence Avenue	e, SW, Washington, D.C. 20250-9410 or call (202)			

200500325

GROUP GERMPLASM ASSIGNMENT

VTIP 03.017

VA97W-24

VTIP 03.018

VA98W-586

VTIP 03:019

VA98W-706#

VTIP 03.020

VA98W-750

VTIP 03.021

VA98W-817

VTIP 03.022

DAN/VA97B-388 Barley

VTIP 03.023

DOYCE/VA00H-137 Barley

VTIP 03.024

VT67 and VT120 Soybean

VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY (hereinafter referred to as the "UNIVERSITY"), assigns to VIRGINIA TECH INTELLECTUAL PROPERTIES, INC. (hereinafter referred to as "VTIP") all rights, title and interest in and to all of the above-listed GERMPLASMS as held by the UNIVERSITY.

The UNIVERSITY, by its authorized agents, agrees that it will execute all necessary assignments as requested by VTIP, to facilitate the filing of patent applications and/or copyright registrations. It will render any reasonable assistance requested to aid in preparation of such applications and/or registrations.

The UNIVERSITY shall retain the right to make use of the GERMPLASMS for internal research and other non-commercial purposes without cost to the UNIVERSITY.

All royalties, rents, payments, or any cash receipts from the sale, assignment, transfer, licensing or use of the GERMPLASMS shall be the property of VTIP and shall be distributed according to the provisions of the Virginia Agricultural Experiment Station (VAES) Plant Germplasm Release Policy (PGRP).

Prior to the execution of this Assignment, the UNIVERSITY has not granted the right of license to make, use, or sell said GERMPLASMS to anyone except to VTIP, nor has it otherwise encumbered its rights, title and interest in said GERMPLASMS, and it will not execute any instrument in conflict with this Assignment.

IN	WITNESS	WHEREOF,	the UNIVERSITY h	as caused thi	s Assignment to be
signed this	25	day of	the UNIVERSITY I		, 2003.

VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY

MINNIS E. RIDENOUR

Chief Operating Officer

STATE OF VIRGINIA

COUNTY OF MONTGOMERY, to-wit:

The foregoing instrum	ent was acknowle	dged before m	e this <u>25</u> day of
March	, 2003, by_	Mixuo	& Ridensur
of Virginia Polytechnic Institu	te and State Unive	ersity, on behal	If of said University.
	11	11	
Notary Public	Gerry M.	Chenau	et
Notary Public	0		
My commission	expires 2	130/17	7.